Diabetes Research and Clinical Practice, 8 (1990) 289–290
Elsevier

Key Word Index to Volume 8 (1990)

Albumin
– Glycation; Clearance; Proteinuria, 221
Asahikawa colony (CHAD)
– Diabetic Chinese hamster; Immunoperoxidase method; Morphometric analysis; Islet cell composition, 201
Autoantibody
– Insulin; Hypoglycemia; Carbimazole, 169

Carbimazole
– Autoantibody; Insulin; Hypoglycemia, 169
Children, diabetic
– Exocrine pancreatic function; PABA; Trypsinemia test, 263
Clearance
– Glycation; Albumin; Proteinuria, 221

Diabetic Chinese hamster
– Asahikawa colony (CHAD); Immunoperoxidase method; Morphometric analysis; Islet cell composition, 201
Diabetic neuropathy
– Erythrocyte; Na, K-ATPase (sodium pump); Urinary C-peptide immunoreactivity, 177
Diabetic rats
– Estrogen; Insulin-like growth factor I; Insulin treatment, 235
Diabetogenesis
– Offspring of conjugal diabetics; Predictive parameter; Immunoreactive insulin, 269
Diagnostic test
– Fructosamine; Fasting plasma glucose; Hemoglobin A1c; Discriminant analysis, 187
Discriminant analysis
– Fructosamine; Fasting plasma glucose; Hemoglobin A1c; Diagnostic test, 187

Erythrocyte
– Na,K-ATPase (sodium pump); Diabetic neuropathy; Urinary C-peptide immunoreactivity, 177
Estrogen
– Diabetic rats; Insulin-like growth factor I; Insulin treatment, 235
Exocrine pancreatic function
– PABA; Trypsinemia test; Children, diabetic, 263

Fasting plasma glucose
– Fructosamine; Hemoglobin A1c; Diagnostic test; Discriminant analysis, 187
Fetal
– Proislets; Isograft; Mouse, 275
Fructosamine
– Fasting plasma glucose; Hemoglobin A1c; Diagnostic test; Discriminant analysis, 187

Glycated protein
– Immunohistochemistry; Kidney; Proximal tubule, 215
Glycation
– Albumin; Clearance; Proteinuria, 221
Glyoxalase
– HLA; Immunoglobulin heavy-chain allotype (Gm); Properdin factor B(BF), 253

HLA
– Immunoglobulin heavy-chain allotype (Gm); Properdin factor B(BF); Glyoxalase, 253
Hemoglobulin A1c
– Fructosamine; Fasting plasma glucose; Diagnostic test; Discriminant analysis, 187
Hypervariable region
– Insulin gene; Mutation; Tandem repeating oligonucleotide; Origin, 183
Hypoglycemia
– Autoantibody; Insulin; Carbimazole, 169

Immunoglobulin heavy-chain allotype (Gm)
– HLA; Properdin factor B(BF); Glyoxalase, 253
Immunohistochemistry
– Glycated protein; Kidney; Proximal tubule, 215
Immunoperoxidase method
– Diabetic Chinese hamster; Asahikawa colony (CHAD); Morphometric analysis; Islet cell composition, 201
Immunoreactive insulin
– Offspring of conjugal diabetics; Predictive parameter; Diabetogenesis, 269
Impaired glucose tolerance test
– Non-insulin-dependent diabetes mellitus; Prevalence of diabetes; Obesity; Migrants, 227
Insulin
– Autoantibody; Hypoglycemia; Carbimazole, 169
Insulin
– NIDDM; Secondary failure; Oral antidiabetic agent; Placebo, 243
Insulin gene
– Mutation; Hypervariable region; Tandem repeating oligonucleotide; Origin, 183

Insulin release
– Isolated islets; beta-Bungarotoxin; Parasympathetic nerve; Nicotonic receptor, 193
Insulin treatment
– Diabetic rats; Estrogen; Insulin-like growth factor I, 235
Insulin-like growth factor I
– Diabetic rats; Estrogen; Insulin treatment, 235
Islet cell composition
– Diabetic Chinese hamster; Asahikawa colony (CHAD); Immunoperoxidase method; Morphometric analysis, 201
Isograft
– Fetal; Proislets; Mouse, 275
Isolated islets
– Insulin release; beta-Bungarotoxin; Parasympathetic nerve; Nicotonic receptor, 193

Kidney
– Immunohistochemistry; Glycated protein; Proximal tubule, 215

Microangiopathy
– Nephropathy; Retinopathy, 283
Migrants
– Non-insulin-dependent diabetes mellitus; Impaired glucoase tolerance test; Prevalence of diabetes; Obesity, 227
Morphometric analysis
– Diabetic Chinese hamster; Asahikawa colony (CHAD); Immunoperoxidase method; Islet cell composition, 201
Mouse
– Fetal; Proislets; Isograft, 275
Mutation
– Insulin gene; Hypervariable region; Tandem repeating oligonucleotide; Origin, 183

NIDDM
– Secondary failure; Oral antidiabetic agent; Insulin; Placebo, 243
Na, K-ATPase (sodium pump)
– Erythrocyte; Diabetic neuropathy; Urinary C-peptide immunoreactivity, 177
Nephropathy
– Retinopathy; Microangiopathy, 283
Nicotonic receptor
– Insulin release; Isolated islets; beta-Bungarotoxin; Parasympathetic nerve, 193
Non-insulin-dependent diabetes mellitus
– Impaired glucose tolerance test; Prevalence of diabetes; Obesity; Migrants, 227

Obesity
– Non-insulin-dependent diabetes mellitus; Impaired glucose tolerance test; Prevalence of diabetes; Migrants, 227
Offspring of conjugal diabetics
– Predictive parameter; Immunoreactive insulin; Diabetogenesis, 269
Oral antidiabetic agent
– NIDDM; Secondary failure; Insulin; Placebo, 243
Origin
– Insulin gene; Mutation; Hypervariable region; Tandem repeating oligonucleotide, 183

PABA
– Exocrine pancreatic function; Trypsinemia test; Children, diabetic, 263
Parasympathetic nerve
– Insulin release; Isolated islets; beta-Bungarotoxin; Nicotonic receptor, 193
Placebo
– NIDDM; Secondary failure; Oral antidiabetic agent; Insulin, 243
Predictive parameter
– Offspring of conjugal diabetics; Immunoreactive insulin; Diabetogenesis, 269
Prevalence of diabetes
– Non-insulin-dependent diabetes mellitus; Impaired glucose tolerance test; Obesity; Migrants, 227
Proislets
– Fetal; Isograft; Mouse, 275
Properdin factor B(BF)
– HLA; Immunoglobulin heavy-chain allotype (Gm); Glyoxalase, 253
Proteinuria
– Glycation; Albumin; Clearance, 221
Proximal tubule
– Immunohistochemistry; Glycated protein; Kidney, 215

Retinopathy
– Nephropathy; Microangiopathy, 283

Secondary failure
– NIDDM; Oral antidiabetic agent; Insulin; Placebo, 243

Tandem
– Insulin gene; Mutation; Hypervariable region; Origin, 183
Trypsinemia test
– Exocrine pancreatic function; PABA; Children, diabetic, 263

Urinary
– Erythrocyte; Na, K-ATPase (sodium pump); Diabetic neuropathy, 177

beta-Bungarotoxin
– Insulin release; Isolated islets;: Parasympathetic nerve; Nicotonic receptor, 193

Diabetes Research and Clinical Practice, 8 (1990) 291–292
Elsevier

Author Index to Volume 8 (1990)

Abdella, N., Salman, A. and Moro, M., Classical microangiopathic diabetic complications in the absence of overt diabetes mellitus, 283
Alberti, K.G.M.M., see Swai, A.B.M., 227
Allen, T.J., see Layton, G.J., 221
Aono, S., see Mimura, G., 253
Armitage, M., Franklyn, J., Scott-Morgan, L., Parr, J., Borsey, D.Q., Sheppard, M. and Wilkin, T.J., Insulin autoantibodies in Graves' disease – before and after carbimazole therapy, 169
Awata, T., Iwamoto, Y., Matsuda, A. and Kuzuya, T., Hypervariable region 5′-flanking [Leu A3]insulin gene of insulin Tochigi is different from those of insulin Wakayama I,II, 183

Baba, S., see Ejiri, K., 193
Borsey, D.Q., see Armitage, M., 169

Chuwa, L.M., see Swai, A.B.M., 227
Cortona, L., see Lorini, R., 263

Doi, K., see Mimura, G., 253

Ejiri, K., Taniguchi, H., Ishihara, K., Hara, Y. and Baba, S., Possible involvement of cholinergic nicotinic receptor in insulin release from isolated rat islets, 193
Elkind-Hirsch, K.E., see Rogers, D.G., 235

Franklyn, J., see Armitage, M., 169

Hara, Y., see Ejiri, K., 193
Hashimoto, T., see Noda, K., 177
Hibi, I., see Mimura, G., 253
Huupponen, R., see Sotaniemi, E.A., 243

Ikeda, Y., see Mimura, G., 253
Ishihara, K., see Ejiri, K., 193
Iwamoto, Y., see Awata, T., 183
Iwashima, Y., Watanabe, K. and Makino, I., Changes in the pancreatic A-, B- and D-cell populations during development of diabetes in spontaneously diabetic Chinese hamsters of the Asahikawa colony (CHAD), 201

Jerums, G., see Layton, G.J., 221
Jinnouchi, T., see Mimura, G., 253

Karvonen, I., see Sotaniemi, E.A., 243
Kermali, W., see Swai, A.B.M., 227
Kida, K., see Mimura, G., 253
Kitagawa, T., see Mimura, G., 253
Kobayashi, T., see Mimura, G., 253
Kuzuya, H., see Mimura, G., 253
Kuzuya, T., see Awata, T., 183

Layton, G.J., Allen, T.J. and Jerums, G., Longitudinal evaluation of the renal clearance of glycated albumin in the diabetic rat, 221
Lorini, R., Cortona, L., Scotta, M.S., Melzi d'Eril, G.V. and Severi, F., Exocrine pancreatic function in children and adolescents with insulin-dependent diabetes mellitus, 263
Lukmanji, Z., see Swai, A.B.M., 227

Mackie, D.J., see Simeonovic, C.J., 275
Makene, W., see Swai, A.B.M., 227
Makino, I., see Iwashima, Y., 201
Maro, E., see Swai, A.B.M., 227
Matsudo, A., see Awata, T., 183
Matsuura, N., see Mimura, G., 253
McLarty, D.G., see Swai, A.B.M., 227
Melzi d'Eril, G.V., see Lorini, R., 263
Nimura, G., Kida, K., Matsuura, N., Toyota, T., Kitagawa, T., Kobayashi, T., Hibi, I., Ikeda, Y., Tuchida, I., Kuzuya, H., Aono, S., Doi, K., Nishimukai, H., Jinnouchi, T. and Murakami, K., Immunogenetics of early-onset insulin-dependent diabetes mellitus among the Japanese: HLA, Gm, BF, GLO, and organ-specific autoantibodies – the J.D.S. study, 253
Miyazawa, K., see Sekikawa, A., 187
Mizuno, A., see Yano, M., 215
Mohan, V., see Ramachandran, A., 269
Moro, M., see Abdella, N., 283
Murakami, K., see Mimura, G., 253

Nawata, H., see Noda, K., 177
Nishimukai, H., see Mimura, G., 253

Noda, K., Umeda, F., Hashimoto, T., Yamashita, T. and Nawata, H., Erythrocytes from diabetes with neuropathy have fewer sodium pumps, 177

Parr, J., see Armitage, M., 169

Ramachandran, A., Snehalatha, C., Mohan, V. and Viswanathan, M., Development of carbohydrate intolerance in offspring of Asian Indian conjugal type 2 diabetic parents, 269
Rogers, D.G., Valdes, C.T. and Elkind-Hirsch, K.E., The effect of ovarian function on insulin-like growth factor I plasma levels and hepatic IGF-I mRNA levels in diabetic rats treated with insulin, 235
Rytömaa, K., see Sotaniemi, E.A., 243

Salman, A., see Abdella, N., 283
Sasaki, H., see Sekikawa, A., 187
Scott-Morgan, L., see Armitage, M., 169
Scotta, M.S., see Lorini, R., 263
Sekikawa, A., Tominaga, M., Takahashi, K., Watanabe, H., Miyazawa, K. and Sasaki, H., Is examination of fructosamine levels valuable as a diagnostic test for diabetes mellitus?, 187
Severi, F., see Lorini, R., 263
Sherrif, F., see Swai, O.B.M., 227
Sheppard, M., see Armitage, M., 169
Shima, K., see Yano, M., 215
Simeonovic, C.J., Mackie, D.J. and Wilson, J.D., Functional proporties of proislet isografts in diabetic mice, 275
Snehalatha, C., see Ramachandran, A., 269
Sotaniemi, E.A., Vierimaa, E., Huupponen, R., Karvonen, I. and Rytömaa, K., Insulin and sulphonylurea in the therapy of type 2 diabetes, 243
Swai, A.B.M., McLarty, F., Sherrif, D.G., Chuwa, L.M., Maro, E., Lukmanji, Z., Kermali, W., Makene, W. and Alberti, K.G.M.M., Diabetes and impaired glucose tolerance in an Asian community in Tanzania, 227

Takahashi, K., see Sekikawa, A., 187
Taniguchi, H., see Ejiri, K., 193
Tominaga, M., see Sekikawa, A., 187
Toyota, T., see Mimura, G., 253
Tuchida, I., see Mimura, G., 253

Umeda, F., see Noda, K., 177

Valdes, C.T., see Rogers, D.G., 235
Vierimaa, E., see Sotaniemi, E.A., 243
Viswanathan, M., see Ramachandran, A., 269

Watanabe, H., see Sekikawa, A., 187
Watanabe, K., see Iwashima, Y., 201
Wilkin, T.J., see Armitage, M., 169
Wilson, J.D., see Simeonovic, C.J., 275

Yamashita, T., see Noda, K., 177
Yano, M., Mizuno, A. and Shima, K., Immunohistochemical localization of glycated protein in diabetic rat kidney, 215